

p-JNK (G-7): sc-6254



The Power to Question

BACKGROUND

The mitogen-activated protein (MAP) kinases ERK-1 and ERK-2 are proline-directed kinases that are activated through concomitant phosphorylation of tyrosine and threonine residues. The JNK family, which includes JNK1, JNK2 and JNK3, is distantly related to the MAP kinase family, members of which are activated by dual phosphorylation at a Thr-Pro-Tyr motif, specifically at Thr 183 and Tyr 185 residues, in response to ultraviolet (UV) light. This motif is divergent from the Thr-Glu-Tyr motif characteristic of the MAP kinase family. JNK is phosphorylated by JNK-activating kinase (JNKK1 and JNKK2), which are members of the MEK family. Activated JNK mediates the phosphorylation of c-Jun at the amino-terminal serine regulatory sites, Ser 63 and Ser 73, which stimulates the transactivation function of c-Jun.

SOURCE

p-JNK (G-7) is a mouse monoclonal antibody raised against a sequence containing Thr 183 and Tyr 185 phosphorylated JNK of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

p-JNK (G-7) is available conjugated to agarose (sc-6254 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-6254 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-6254 PE), fluorescein (sc-6254 FITC), Alexa Fluor® 488 (sc-6254 AF488), Alexa Fluor® 546 (sc-6254 AF546), Alexa Fluor® 594 (sc-6254 AF594) or Alexa Fluor® 647 (sc-6254 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-6254 AF680) or Alexa Fluor® 790 (sc-6254 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

In addition, p-JNK (G-7) is available conjugated to biotin (sc-6254 B), 200 µg/ml, for WB, IHC(P) and ELISA; and to either TRITC (sc-6254 TRITC), 200 µg/ml or Alexa Fluor® 405 (sc-6254 AF405), 100 µg/2 ml, for IF, IHC(P) and FCM.

Blocking peptide available for competition studies, sc-6254 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

p-JNK (G-7) is recommended for detection of Thr 183 and Tyr 185 phosphorylated JNK1, JNK2 and JNK3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of p-JNK p46 isoform: 46 kDa.


Molecular Weight of p-JNK p54 isoform: 54 kDa.

Positive Controls: NIH/3T3 + anisomycin cell lysate: sc-2247, NIH/3T3 + UV cell lysate: sc-3804 or RAW 264.7 + UV cell lysate: sc-24769.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



Western blot analysis of JNK phosphorylation in untreated (A, C) and anisomycin treated (B, D) Jurkat whole cell lysates. Antibodies tested include p-JNK (G-7): sc-6254 (A, B) and JNK (FL): sc-571 (C, D).

p-JNK (G-7) Alexa Fluor® 488: sc-6254 AF488. Direct immunofluorescence staining of formalin-fixed SW480 cells showing nuclear and cytoplasmic localization. Blocked with UltraCruz® Blocking Reagent: sc-516214 (A). p-JNK (G-7): sc-6254. Immunoperoxidase staining of formalin fixed, paraffin-embedded human breast tissue showing cytoplasmic and membrane staining of glandular cells and myoepithelial cells (B).

SELECT PRODUCT CITATIONS

- Rincon, M., et al. 1998. Interferon- γ expression by Th1 effector T cells mediated by the p38 MAP kinase signaling pathway. *EMBO J.* 17: 2817-2829.
- Yamboliev, I.A., et al. 1998. Endothelin-1 activates MAP kinases and c-Jun in pulmonary artery smooth muscle. *Pulm. Pharmacol. Ther.* 11: 205-208.
- Mukaro, V.R., et al. 2018. Small tumor necrosis factor receptor biologics inhibit the tumor necrosis factor-p38 signalling axis and inflammation. *Nat. Commun.* 9: 1365.
- Gibbs, K.L., et al. 2018. Inhibiting p38 MAPK α rescues axonal retrograde transport defects in a mouse model of ALS. *Cell Death Dis.* 9: 596.
- Gorojod, R.M., et al. 2018. Heme Oxygenase-1 protects astroglia against manganese-induced oxidative injury by regulating mitochondrial quality control. *Toxicol. Lett.* 295: 357-368.
- Yu, T., et al. 2018. Exclusive enteral nutrition protects against inflammatory bowel disease by inhibiting NF κ B activation through regulation of the p38/MSK1 pathway. *Int. J. Mol. Med.* 42: 1305-1316.
- Juan, T.K., et al. 2018. Tetrandrine suppresses adhesion, migration and invasion of human colon cancer SW620 cells via inhibition of nuclear factor- κ B, matrix metalloproteinase-2 and matrix metalloproteinase-9 signaling pathways. *Oncol. Lett.* 15: 7716-7724.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA